

lution as the standard solution. Perform the test with these solutions as directed under the Thin-layer Chromatography. Spot 40 μL each of the sample solution and the standard solution on a plate of silica gel with fluorescent indicator for thin-layer chromatography. Develop the plate with a mixture of chloroform, methanol, acetone and ammonia solution (28) (30:10:10:1) to a distance of about 12 cm, and air-dry the plate. Examine under ultraviolet light (main wavelength: 254 nm): the spots other than the principal spot from the sample solution are not more intense than the spot from the standard solution.

Loss on drying Not more than 0.20% (1 g, in vacuum, 105°C, 4 hours).

Residue on ignition Not more than 0.10% (1 g).

Assay Weigh accurately about 1.5 g of Arotinolol Hydrochloride, previously dried, dissolve in dimethylsulfoxide to make exactly 25 mL. Pipet 5 mL of this solution, add 100 mL of water and 5 mL of sodium hydroxide TS, and extract with three 50-mL portions of dichloromethane. Filter each dichloromethane extract through a pledget of absorbent cotton with anhydrous sodium sulfate on it. Evaporate combined filtrate to dryness in vacuum. Dissolve the residue in 70 mL of acetic acid (100), and titrate with 0.05 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.

Each mL of 0.05 mol/L perchloric acid VS
= 20.400 mg of $\text{C}_{15}\text{H}_{21}\text{N}_3\text{O}_2\text{S}_3\cdot\text{HCl}$

Containers and storage Containers—Tight containers.
Storage—Light-resistant.

Arsenic Trioxide

Arsenous Acid

三酸化ヒ素

As_2O_3 : 197.84

Arsenic Trioxide, when dried, contains not less than 99.5% of As_2O_3 .

Description Arsenic Trioxide occurs as a white powder.

It is odorless. It is practically insoluble in water, in ethanol (95) and in diethyl ether.

It dissolves in sodium hydroxide TS.

Identification Dissolve 0.2 g of Arsenic Trioxide in 40 mL of water by heating on a water bath: the solution responds to the Qualitative Tests for arsenite.

Purity Clarity of solution—To 1.0 g of Arsenic Trioxide add 10 mL of ammonia TS, and heat gently: the solution is clear.

Loss on drying Not more than 0.5% (1 g, 105°C, 3 hours).

Assay Weigh accurately about 0.15 g of Arsenic Trioxide, previously dried, dissolve in 20 mL of a solution of sodium hydroxide (1 in 25), by warming, if necessary. Add 40 mL of water and 2 drops of methyl orange TS, then add dilute

hydrochloric acid until the color of the solution becomes light red. Add 2 g of sodium hydrogen carbonate and 50 mL of water to this solution, and titrate with 0.05 mol/L iodine VS (indicator: 3 mL of starch TS).


Each mL of 0.05 mol/L iodine VS = 4.946 mg of As_2O_3

Containers and storage Containers—Tight containers.

Ascorbic Acid

Vitamin C

アスコルビン酸



$\text{C}_6\text{H}_8\text{O}_6$: 176.12

2,3-Didehydro-L-threo-hexono-1,4-lactone [50-81-7]

Ascorbic Acid, when dried, contains not less than 99.0% of L-ascorbic acid ($\text{C}_6\text{H}_8\text{O}_6$).

Description Ascorbic Acid occurs as white crystals or a white, crystalline powder. It is odorless, and has an acid taste.

It is freely soluble in water, sparingly soluble in ethanol (95), and practically insoluble in diethyl ether.

Melting point: about 190°C (with decomposition).

Identification (1) To 5 mL each of a solution of Ascorbic Acid (1 in 50) add 1 drop of potassium permanganate TS or 1 to 2 drops of 2,6-dichloroindophenol sodium TS: the color of the solution is discharged immediately in each case.

(2) Dissolve 0.1 g of Ascorbic Acid in 100 mL of a solution of metaphosphoric acid (1 in 50). To 5 mL of the solution add iodine TS until the color of the solution becomes light yellow. Then add 1 drop of a solution of copper (II) sulfate pentahydrate (1 in 1000) and 1 drop of pyrrole, and warm the mixture at 50°C for 5 minutes: a blue color develops.

Optical rotation $[\alpha]_D^{20}$: + 20.5 – + 21.5° (2.5 g, water, 25 mL, 100 mm).

pH Dissolve 1.0 g of Ascorbic Acid in 20 mL of water: the pH of this solution is between 2.2 and 2.5.

Purity (1) Clarity and color of solution—Dissolve 1.0 g of Ascorbic Acid in 20 mL of water: the solution is clear and colorless.

(2) Heavy metals—Perform the test with 1.0 g of Ascorbic Acid according to Method 1. Prepare the control solution with 2.0 mL of Standard Lead Solution (not more than 20 ppm).

Loss on drying Not more than 0.20% (1 g, silica gel, 24 hours).

Residue on ignition Not more than 0.10% (1 g).

Assay Weigh accurately about 0.2 g of Ascorbic Acid, previously dried, dissolve in 50 mL of a solution of